



Angolan cones Species List

Based on WORMS 2018/2019.

The majority of Angolan cones are in the subgenus *Conus (Lautoconus)* Monterosato, 1923.

The exceptions are *C. genuanus* Linnaeus, 1758 and *C. pulcher* [Lightfoot, 1786] in *Conus (Kalloconus)*, *C. ermineus* Born, 1778 in *Conus (Chelyconus)*, *C. tabidus* Reeve, 1844 in *Conus (Monteiroconus)* and *Conus carnalis* Sowerby iii, 1879 in *Conus (Pseudonoduloconus)*.

C. genuanus Linnaeus, 1758; Luanda to Namibe. Senegal to Angola
C. tabidus Reeve, 1844. CVI, Senegal to Angola
C. pulcher [Lightfoot, 1786] Senegal to Angola
C. ermineus Born, 1778 Senegal to Angola

C. aemulus Reeve, 1844; Endemic to Luanda Province, Angola
C. xicoi Röckel, 1987; Endemic to Luanda Province, Angola

Species(32) found only in Benguela and Namibe provinces, Angola. All except *C. carnalis*, in *Lautoconus*.

C. africonus Kiener, 1848.
C. albuquerquei Trovão, 1978.
C. alexandrinus Kaicher, 1977. Synonyms include *C. musivus* Trovão, 1975, *C. tevesi* Trovão, 1978.
C. allaryi Bozzetti, 2008.
C. anabelae Rolán & Röckel, 2001.
C. annegretae Schönherr, 2018.
C. babaensis Rolán & Röckel, 2001.
C. bocagei Trovão, 1978.
C. bulbosus Reeve, 1843.
C. carnalis Sowerby III, 1879.
C. cepasi Trovão, 1975.
C. chytreus Tryon, 1884.
C. equiminaensis Schönherr, 2018.
C. eusebioi Schönherr, 2018.
C. filmeri Rolán & Röckel, 2000.
C. flavusalbus Rolán & Röckel, 2000.
C. franciscoi Rolán & Röckel, 2000.
C. fuscolineatus Sowerby III, 1905.
C. inesa Monteiro A., Afonso C., Tenorio M.J., Rosado J. & Pirinhas D. 2014.
C. lineopunctatus Kaicher, 1977. Synonyms include *C. neoguttatus* Da Motta, 1991.
C. lobitensis Kaicher, 1977.
C. medvedevi Monteiro A., Afonso C., Tenorio M.J., Rosado J. & Pirinhas D. 2014.

C. micropunctatus Rolán & Röckel, 2000.
C. naranjus Trovão, 1975.
C. negroides Kaicher, 1977. Synonyms include *C. gabriellae* Rolán & Röckel, 2000.
C. nobrei Trovão, 1975.
C. nunesi Schönherr, 2018.
C. petuchi Monteiro A., Afonso C., Tenorio M.J., Rosado J. & Pirinhas D. 2014.
C. tenuilineatus Rolán & Röckel, 2001.
C. trovai Rolán & Röckel, 2000.
C. variegatus Kiener, 1848.
C. zebroides Kiener, 1848.

C. obtusus Kiener, 1850 considered a synonym of *C. variegatus*. (often used as a form name).
C. dealbatus Adams, 1854 considered a nomen dubium, possibly a fossil species (name often used for white shells).

Page last updated 9 May 2019.